
1 Introduction

J.D. HANSEN¹ AND J.A. JOHNSON²

¹USDA-ARS Yakima Agricultural Research Laboratory, Wapato, Washington, USA; e-mail: jimbob@yarl.ars.usda.gov ; ²USDA-ARS San Joaquin Valley Agricultural Sciences Center, Parlier, California, USA; e-mail: jjohnson@fresno.ars.usda.gov

1.1 History and Purpose of Quarantine and Phytosanitation Requirements

Introduction

Heat has had a variety of uses since primitive times, such as in cooking and food preservation, but its use was limited as a pest control method for stored products until the modern era. Heat can be generated by various ways: chemical oxidation, combustion, electrical resistance and electromagnetic exposure, and heat treatments have been devised that take advantage of each.

The manner in which heat is produced affects both products and their pests, and the success of a given treatment depends on its ability to control insects without causing product damage. This chapter will briefly review the history and development of product treatments in general and heat treatments in particular that meet pest phytosanitation and quarantine security requirements. Terms used in the regulatory processes will be defined, and a short review of the current status of heat treatments will be presented.

Treatments

Postharvest treatments are a recent development. For most of human history, insect pests in stored products have been tolerated. However, two events led to the development of postharvest methods to eliminate insects. The first relates to the distribution of pest insects, which historically was limited by their biology and geophysical forces. With increased travel by humans – particularly wide-ranging exploration and commerce – insects can now be transported into new areas. If the insect has potential to do damage, even though it is a minor pest in its native range, measures that prevent its introduction and establishment are

needed. This concern has resulted in a series of procedures and regulations pertaining to quarantine insects.

The second event that led to the development of postharvest methods to eliminate insects involved the rapid evolution of agricultural technology to store surplus commodities until future consumption or transport to distant markets for greater economic gain (see Table 1.1, which illustrates the huge global market for two particular commodities, fruit and nuts). To prevent damage during storage by various insects – particularly beetles and moths, phytosanitation procedures were developed to reduce and control the populations of these pests. Because of the complexities of modern commerce, the distinction between quarantine and phytosanitation is often unclear.

Quarantine Regulations

Quarantine statutes involve the treatment or shipment of an agricultural commodity from one location to another. They can be imposed by various bodies: (i) local governmental jurisdiction, such as the Washington State (USA) quarantine against intrastate movement of apples from areas where fruits have been infested with the apple maggot, *Rhagoletis pomonella* (Walsh) (Diptera:

Table 1.1. World export of fruit and nut products.

Commodity	Four-year average (2000–2003)	
	Quantity (metric tonnes)	Value (US\$1,000)
<i>Fruits</i>		
Cherries	155,769	388,011
Dates	494,863	260,179
Figs (fresh and dried)	88,270	141,338
Grapefruit and Pomelos	1,037,373	475,102
Grapes	2,806,023	2,701,020
Lemons and limes	1,737,864	796,061
Mangoes	714,655	437,758
Olives	36,813	34,761
Oranges	4,783,691	2,024,224
Papayas	209,718	132,112
Peaches and Nectarines	1,222,043	1,007,580
Pineapples	1,294,431	589,146
Plums	434,231	351,183
Prunes (dried plums)	160,846	267,899
Raisins	632,459	607,248
Strawberries	492,654	789,275
Tangerines	2,564,985	1,559,091
<i>Nuts (shelled and in-shell)</i>		
Almonds	393,268	1,099,125
Hazelnuts	224,843	557,993
Pistachios	211,212	697,965
Walnuts	186,885	460,667

Tephritidae); (ii) within national boundaries, such as the quarantine against movement of cherries from the US Pacific Northwest states to California (CDFA, 1998) because of possible infestation by cherry fruit fly, *Rhagoletis cingulata* (Loew) (Diptera: Tephritidae); or (iii) between nations, such as the quarantine of possible host materials of the codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae), in exports to Japan (MAFF-Japan, 1950).

To facilitate the implementation of such quarantines, groups of nations within a region have formed organizations that provide similar regulations against a common potential pest. Laws involving the importation of commodities that may contain the Caribbean fruit fly, *Anastrepha suspense* (Loew) (Diptera: Tephritidae), imposed by nations in the European and Mediterranean Plant Protection Organization (EPPO), are good examples.

Other international regulatory organizations include the Caribbean Plant Protection Commission, Inter-African Phytosanitary Council and the North American Plant Protection Organization. Individual countries have their own plant protection agencies, such as the Australian Quarantine Inspection Service, Ministry of Agriculture, Forestry and Fisheries (Japan) and Animal and Plant Health Inspection Service (USA). Regulation within a country often falls on a local or state government body, such as the US California Department of Food and Agriculture and Washington State Department of Agriculture.

All of the above entities may write their own regulations to meet immediate concerns and generate lists of quarantine pests. For example, the EPPO has a list for quarantine pests not present (A1 pests) and another for quarantine pests present but not widely distributed and officially under control (A2 pests). The state of California has a similar rating system, with 'A' pests requiring that infested commodities be rejected or treated, and 'Q' pests temporarily requiring the same action as A pests until a permanent rating can be determined. To assure exclusion of the quarantine pest, these lists include all potential hosts.

Enforcement of quarantine regulations relies primarily on international trade agreements and standards that may include the use of specific treatments on select commodities to assure quarantine security. Hence, international commerce in commodities potentially containing quarantine pests is strictly regulated.

Phytosanitation

Phytosanitation generally does not involve regulation as extensive as that applied to quarantine. Rather, economic considerations direct its application. It benefits the producer to promote commodities free of pests, both to restrain internal costs and to promote an attractive product. The burden of phytosanitation usually falls on the fruit or produce packer or marketer, and treatments may be applied repeatedly.

Probit-9 and Quarantine Security

The standard of quarantine security for many importing countries is probit-9, or 99.996832% mortality of the pest population (Robertson *et al.*, 1994b). In practical terms, probit-9 means that only 32 individuals can survive out of 1 million treated pests. Originally suggested by Baker (1939), the use of probit-9 was directed at those not familiar with sophisticated experimentation and data analysis. Baker used moisture-saturated hot air to control the Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae), on Hawaiian kamani nuts shipped to the American mainland, and intended that the criteria used to calculate probit-9 mortality be successful adult emergence from a single batch of insects.

Probit-9 security levels are normally applied to quarantine treatments of commodities that have a high and frequently unknown level of pest infestation. In practice, this condition is often ignored. For example, cherries imported to Japan from the Pacific Northwest of the USA must be treated with quarantine procedures that have demonstrated probit-9 efficacy on the last codling moth instar, even though intense postharvest inspections demonstrate its rare occurrence. Between 1978 and 1996, one potential codling moth larva was identified from out of 4.9×10^8 individually inspected fruits from the Pacific Northwest region of the USA (Wearing *et al.*, 2001).

In developing a new quarantine procedure, it is very difficult to produce the probit-9 level by treating 1 million insects at one time. Instead, a probit model is developed based on a dose-response curve from much smaller populations, and then projected out to the probit-9 level to determine the quarantine treatment. A statistical test may be included to determine whether the original data fit a probit distribution. Confirmatory tests are then performed, during which the proposed quarantine treatment is tested against a large number of pest insects.

Some countries require a specific number of test insects to be treated during confirmatory studies: 30,000 test insects with no survivors are commonly used. Although this approach is flawed by several severe statistical problems, described in detail elsewhere (Landolt *et al.*, 1984; Chew, 1994; Robertson *et al.*, 1994a, b), many regulatory bodies continue to base their quarantine programs on probit-9 security. A more thorough discussion of the mathematical approaches for determining efficacy is given in Chapter 5, this volume.

1.2 Review of Treatments

General Overview

Quarantine and phytosanitation treatments can be sorted into several categories: chemical, biological, irradiation and physical. Briefly, chemical approaches include a wide assortment of distinct compounds: (i) fumigants

(hydrogen cyanide, phosphine, carbon disulphide, halogenated hydrocarbons such as ethylene dibromide and methyl bromide, and acrylonitrile); (ii) dips (insecticides and soaps); and (iii) controlled or modified atmospheres (often used in combination with a temperature treatment).

Fumigants are the most prevalent, but are losing support because of health, environmental and safety concerns (Stark, 1994; Yokoyama, 1994). The postharvest alternatives to methyl bromide fumigation will be discussed later in this chapter. Controlled atmospheres (CA), reviewed by Carpenter and Potter (1994) and Hallman (1994), involve changing the composition of gases, usually substituting oxygen with nitrogen or carbon dioxide. Insecticides used in dips and sprays include soaps, organophosphorus compounds, organochlorines and insect growth regulators, and these have been reviewed by Hansen and Hara (1994) and Heather (1994). Approval of chemical approaches varies between different jurisdictions.

Biological approaches include the use of natural enemies or microbial agents (generally limited to phytosanitary treatments of stored grains or other durables), host-plant resistance (particularly among cultivars) and pest-free zones, which may be geographical or temporal. For pest-free zones to be effective, the host-pest relationship must be well known, and site manipulation – such as the removal of alternative hosts as well as intensive sampling – may be required. Armstrong (1994a), Greany (1994), Riherd *et al.* (1994), Moore *et al.* (2000) and Schöller and Flinn (2000) have all reviewed different biological methods.

Irradiation, including exposure to gamma and X-rays, is an old technology where dose mortality schedules have been established for several pests on a variety of commodities. Safety, expense, environmental concerns and consumer acceptance interfere with its widespread use. For further information, consult Moy (1985), Burditt (1994) and Nation and Burditt (1994).

Physical methods of quarantine and phytosanitation treatments include mechanical, ultrasound, vacuum and temperature extremes. Mechanical systems use brushes and water sprays (Walker *et al.*, 1996; Whiting *et al.*, 1998; Prusky *et al.*, 1999). Ultrasound produces alternating high and low pressure waves that cause cavitation at the cellular level (Sala *et al.*, 1999) to reduce surface pests on fruit (Hansen, 2001).

Vacuum treatments, applied primarily to durable commodities, are gaining interest due to the development of flexible, low-cost treatment containers (Navarro *et al.*, 2001). Alone, vacuum treatments appear to cause mortality in insects by creating a low-oxygen atmosphere and, in low-moisture environments, may cause rapid dehydration (Navarro and Calderon, 1979). A vacuum microwave (MW) grain dryer was found to rapidly disinfest stored grain of insects (Tilton and Vardell, 1982a, b).

Cold temperature treatments, particularly cold storage, have practical application for many commercial operations (Armstrong, 1994b; Gould, 1994), such as holding commodities before marketing – as well as pest control. Thermal treatments must be precise because of the narrow margin between efficacy and commodity tolerance to the temperature, particularly in fresh fruit and vegetables. Heat treatments have been accepted for commodities entering the USA (USDA, 2005a) and for interstate shipments (see Table 1.2) (USDA,

Table 1.2. Quarantine heat treatments in the USA (from USDA, 2005b, c).

Commodity	Target pests	Treatment schedule
<i>Hot water immersion</i>		
Limes	Mealybugs and other surface pests	49°C or above for 20 min
Longan (lychee from Hawaii)	<i>Ceratitis capitata</i> , <i>Bactrocera dorsalis</i>	49°C or above for 20 min
Mango	<i>Ceratitis capitata</i> , <i>Anastrepha</i> spp., <i>Anastrepha ludens</i>	46°C for 65–110 min, depending on origin, size and shape of fruit
<i>High-temperature forced air</i>		
Citrus from Mexico, infested areas in the USA	<i>Anastrepha</i> spp.	Raise centre of fruit to 44°C over 90 min, hold at 44°C for 100 min
Citrus from Hawaii	<i>Ceratitis capitata</i> , <i>Bactrocera</i> <i>dorsalis</i> , <i>B. cucurbitae</i>	47.2°C (fruit centre) for at least 4 h total treatment time
Mango from Mexico	<i>Anastrepha ludens</i> , <i>A. oblique</i> , <i>A. serpentina</i>	Until seed surface reaches 48°C
Papaya from Chile, Belize and Hawaii	<i>Ceratitis capitata</i> , <i>Bactrocera</i> <i>dorsalis</i> , <i>B. cucurbitae</i>	47.2°C (fruit centre) for at least 4 h total treatment time
Rambutan from Hawaii	<i>Ceratitis capitata</i> , <i>Bactrocera</i> <i>dorsalis</i>	Raise centre of fruit to 47.2°C over 60 min, hold at 47.2°C for 20 min
<i>Vapour heat treatments</i>		
Bell pepper, aubergine, papaya, pineapple, squash, tomato, courgette from Hawaii	<i>Ceratitis capitata</i> , <i>Bactrocera</i> <i>dorsalis</i> , <i>B. cucurbitae</i>	44.4°C (fruit centre) for 8.75 h (heating rate variable)
Clementine, orange, grapefruit, mango from Mexico	<i>Anastrepha</i> spp.	Raise centre of fruit to 43.3°C over 8 h, hold at 43.3°C for 6 h
Clementine from Mexico (alternate treatment)	<i>Anastrepha</i> spp.	Raise centre of fruit to 43.3°C over 6 h, hold at 43.3°C for 4 h; fruit should be heated rapidly during first 2 h
Lychee from Hawaii	<i>Ceratitis capitata</i> , <i>Bactrocera</i> <i>dorsalis</i>	Raise centre of fruit to 47.2°C over 60 min, hold at 47.2°C for 20 min
Mango from Philippines	<i>Bactrocera</i> spp.	Raise centre of fruit to 46°C over 4 h, hold at 46°C for 10 min
Mango from Taiwan	<i>Bactrocera dorsalis</i>	Raise centre of fruit to 46.5°C, hold at 46.5°C for 30 min
Papaya	<i>Ceratitis capitata</i> , <i>Bactrocera</i> <i>dorsalis</i> , <i>B. cucurbitae</i>	Raise centre of fruit to 47.2°C over 4 h
Yellow pitaya from Colombia	<i>Ceratitis capitata</i> , <i>Anastrepha</i> <i>fraterculus</i>	Raise centre of fruit to 46°C over 4 h, hold at 46°C for 20 min
Rambutan from Hawaii	<i>Ceratitis capitata</i> , <i>Bactrocera</i> <i>dorsalis</i>	Raise centre of fruit to 47.2°C over 60 min, hold at 47.2°C for 20 min

2004). Reviews of heat treatments can be found in Stout and Roth (1983), Armstrong (1994b), Hallman and Armstrong (1994), Sharp (1994a, b) and USDA (2005b). Heat treatments will be discussed in more detail later in this chapter.

Methyl Bromide Fumigation and the Montreal Protocol

For many years, ethylene dibromide was the preferred fumigant against quarantine pests, but was withdrawn in 1984 because of its carcinogenic properties. Since the demise of ethylene dibromide, methyl bromide has become the most popular quarantine treatment (Gaunce *et al.*, 1981; Yokoyama *et al.*, 1990; Moffit *et al.*, 1992). In addition to quarantine uses, methyl bromide has been used extensively for phytosanitary treatments of durables such as dried fruit, nuts, beans, processed foods and pet foods, as well as treatment of processing facilities.

However, methyl bromide was identified by the US Environmental Protection Agency (EPA) under the Federal Clean Air Act (Anon., 1990) and by the Montreal Protocol (Anon., 1995) as having high ozone depletion potential. The EPA mandated the removal of methyl bromide from the chemical register and the phase-out of its production and import into the USA by 31 December 2005.

Similar legislation has occurred in other industrialized countries (see Fig. 1.1). Although methyl bromide fumigation for postharvest quarantine treatments is exempt from the ban, price increases from reduced production, unreliable sources and future restrictions under international agreements (USEPA, 2001) suggest that methyl bromide may become unavailable for quarantine applications as well. Consequently, alternatives to methyl bromide are needed for all postharvest applications.

Practical Alternatives to Methyl Bromide Fumigation

Practical alternatives to methyl bromide must be safe and efficacious, must not reduce product quality, storage life or marketability, must be environmentally acceptable and economically feasible. Some applications require a relatively rapid treatment, either to treat large volumes of product or to meet the needs of specific markets. Although many alternatives have been suggested, most require much longer treatment times, extensive changes to the way processors handle product or substantial capital expenditure.

Alternatives to methyl bromide are easiest to adopt when they serve as drop-in replacements, requiring little change to existing plant infrastructure or




Fig. 1.1. Timeline for the reduction of methyl bromide use globally.

processing procedures. Because other fumigants are considered to be the only such alternatives, they are most likely to be adopted when available. However, issues such as extended treatment times, reduced efficacy and increased costs for some alternative fumigants make them less suitable for certain applications. New fumigants may be developed, but testing and registration of new compounds is a lengthy and expensive process.

Several alternatives have been proposed that do not rely on fumigation for postharvest insect control. These range from single treatments – such as modified atmospheres or vacuum – to combining existing sub-efficacious treatments so that the net efficacy reaches acceptable levels. Related to this is the Systems Approach for quarantine applications, where existing commercial operations, along with treatments that can easily be added without disruption, accumulatively reach the quarantine security level, but not necessarily probit-9. Also, intensive inspection is added to assure compliance.

The industry usually favours this procedure because, with no or few added steps, costs remain low while quality is not adversely affected. This has become the standard for many importing countries, along with a phytosanitation certificate indicating that the commodity has passed the required inspections. Regulatory agencies of both the exporting and importing countries oversee this process. Jang and Moffit (1994) provide a good review of the Systems Approach.

Heat treatments appear as likely candidates for methyl bromide alternatives because: (i) the technology is well established; (ii) they can be applied to a wide range of pests; (iii) there are usually no harmful residues or chemical byproducts; (iv) heat can be generated from a variety of sources; and (v) the costs can usually be controlled. The major impediment is the maintenance of product quality. Dried fruit, nuts and other durables – as well as tropical and subtropical fruit and vegetables – are good candidates for thermal treatments because of their heat tolerances (McDonald and Miller, 1994). However, successful treatments have been developed for temperate fruits as well. Other alternatives to methyl bromide fumigation, in addition to temperature methods, are reviewed by Fields and White (2002) and Vincent *et al.* (2003).

1.3 Survey of Heat Treatments

Early Applications of Heat for Pest Control

Since ancient times, heat in the form of solar energy or fire has been used to control insect pests. Early civilizations killed insect pests in stored grains with the heat of the sun (Cotton, 1963). To control locusts, the Chinese Kingdom of Shang (c.1520–1030 BC) appointed anti-locust officials, who used bonfires to burn collected locusts or repel them (Nevo, 1996). This practice was carried into the 20th century in the USA, when it was suggested for control of chinch bugs (Headlee, 1911) and grasshoppers (Milliken, 1916), a practice that continues to be studied for US rangelands (Vermeire *et al.*, 2004).

In more recent times, the most extensive use of heat treatment has been to control grain insects. For this purpose, heat has been used both as a commodity treatment and as a structural treatment for mills and processing facilities. Thermal treatments against the Angoumois grain moth, *Sitotroga cerealella* (Oliver) (Lepidoptera: Gelechiidae), were used in France for stored grains as early as 1792 (Fields and White, 2002), and later the French used devices known as 'insect mills' for heating infested grain (Dean, 1913).

In the USA as early as 1835, heated rooms were used to control *Sitophilus* spp. (Coleoptera: Curculionidae) in wheat (Oosthuizen, 1935). During the first half of the 20th century, research on heat treatments was conducted on a variety of insects, including (i) the khapra beetle, *Trogoderma granarium* Everts (Coleoptera: Dermestidae) (Husain and Bhasin, 1921); (ii) the red flour beetle, *Tribolium castaneum* (Herbst) F. (Coleoptera: Tenebrionidae) (Grossman, 1931); (iii) the confused flour beetle, *Tribolium confusum* Jacqueline du Val (Coleoptera: Tenebrionidae) (Oosthuizen, 1935); (iv) the Angoumois grain moth (Grossman, 1931; Harukawa, 1941); (v) grain weevils, *Sitophilus* spp. (Back and Cotton, 1924; Grossman, 1931; Tsuchiya, 1943); and (vi) others.

Early work with heat treatments using electromagnetic energy (radio frequency or microwaves) centered primarily on stored-product insects. The lethal thermal effects on insects caused by exposure to radio frequencies were first reported by Lutz (1927) and Headlee and Burdette (1929). Hadjinicolaou (1931) and Whitney (1932) attributed the death of a variety of stored-product pests exposed to high-frequency radio waves to internal heat generated within the body of each insect. Davis (1933) described how radio frequency energy could kill stored-product pests. Mouromtseff (1933) and Ulrey (1936) worked on designing oscillators for use against grain pests. Kuznetzova (1937) showed that the mature larvae of the granary weevil were more resistant to high-frequency electric currents than were other life stages and noted that adults outside the grain were more susceptible than those within.

Heat has been used to disinfest mills and processing plants for some time. Chittenden (1897) recommended using steam on flour mill machinery to control contamination by the Mediterranean flour moth, *Anagast kuehniella* (Zeller) (Lepidoptera: Pyralidae), and recommended 52–60°C for a few hours to kill other grain insects. Dean (1913) noted that the heating of mills in Kansas and other Midwestern states demonstrated the efficacy of this method against all life stages of common mill insects. Pepper and Strand (1935) described how heating the structure of a grain mill to 66°C could control grain pests within 24 h.

Since the late 19th century, heat treatments in the form of hot water dips, vapour heat or hot forced air have been used to treat numerous fresh commodities (Hallman and Armstrong, 1994; Sharp, 1994a). In 1909, one of the earliest attempts at postharvest treatment of a horticultural commodity used immersion of fruit in hot water to control tarsonemid mites (Cohen, 1967).

Vapour heat was first used in Mexico in 1913 to control the Mexican fruit fly, *Anastrepha ludens* (Loew) (Diptera: Tephritidae). Procedures for using vapour heat were also developed to control the Mediterranean fruit fly in Florida citrus (Latta, 1932) and several types of fruits and vegetables in

California (Mackie, 1931). A similar approach was adopted with Texas citrus for fruit fly control (Hawkins, 1932).

Recent Progress in Heat Treatments

Since these early attempts to control insects by thermal methods, the applicable technologies have progressed in mechanical design and theory. Advances in instrumentation now provide accurate measurements of temperature and other treatment variables, and refined techniques have improved precision and replication. The following provides a short review of each of the current thermal methods.

Hot water

Hot ($> 40^{\circ}\text{C}$) water baths and dips are the simplest form of heat treatment. Because of the aqueous medium, treatments are anaerobic, with a rapid energy transfer. Although the procedure has a long history, hot water treatments for quarantine applications have only recently been approved, primarily for fruit flies in tropical and subtropical fruits (Sharp, 1986; Sharp *et al.*, 1988, 1989a, b, c; Sharp and Picho-Martinez, 1990).

Couey *et al.* (1985) and Couey and Hayes (1986) combined ethylene dibromide fumigation with hot water baths to control tephritid fruit flies in Hawaiian papayas. Morgan and Crocker (1986) recommended 49°C water baths for 15 min to control weevils in acorns. For other pests, McLaren *et al.* (1997, 1999) described a commercial-scale 2-min hot water bath (50°C) to control the New Zealand flower thrips, *Thrips obscuratus* (Crawford) (Thysanoptera: Thripidae), on apricots, peaches and nectarines. Other examples of the use of hot water for commercial treatments are found in Chapter 13, this volume, and a discussion of thermal death studies using hot water baths in the laboratory is in Chapter 5, this volume.

Vapour

Like water baths, vapour heat treatments use moisture (as saturated air) to transfer thermal energy, usually involving air movement. As presented earlier in this chapter, vapour heat treatment is an old process. It was developed by Baker (1952) for citrus and is now widely used for papaya, pineapple, bell pepper, eggplant, tomato and zucchini (USDA, 2005c). Sinclair and Lindgren (1955) modified vapour heat treatments for California citrus and avocados, and Seo *et al.* (1974) applied this technology for treatment against the oriental fruit fly, *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae). In Florida, McCoy *et al.* (1994) presented a vapour heat treatment against the eggs of the Fuller rose beetle, *Asynonychus godmani* (Crotch) (Coleoptera: Curculionidae). For more information on vapour heat treatments, see Hallman and Armstrong (1994) and Chapter 13 of this volume.

Forced hot air

Forced hot air treatment is similar to vapour heat treatment, but does not have the moisture component and is a more recent development (Armstrong *et al.*, 1989). Improvements in temperature and moisture monitoring and air delivery have advanced forced hot air treatments (Hallman and Armstrong, 1994). Forced hot air treatments are being devised for commodities normally subjected to vapour heat treatment and have also been applied to new commodities. Their disadvantages are the long treatment durations and sophisticated equipment needed for operation. Also, not all horticultural commodities are suitable, such as avocado (Kerbel *et al.*, 1987).

Heat treatments for durable commodities

Heat has been effective against stored-product pests of durable products such as grains, nuts, dried fruits, wood products and museum artefacts. Most recent work has been carried out on grain pests, and there exists a considerable body of laboratory work on the thermal sensitivity of those (Fields, 1992; Saxena *et al.*, 1992; Mahroof *et al.*, 2003, 2005; Boina and Subramanyam, 2004). Dry heat (82.2°C for 7 min) has been recommended for disinfection of milled products of the khapra beetle, *Trogoderma granarium* Everts (Coleoptera: Dermestidae) – an important quarantine grain pest (Stout and Roth, 1983).

In order to improve the efficiency and reduce the cost, continuous flow, fluidized-bed heating systems for grain have been examined in Australia (Dermott and Evans, 1978; Evans *et al.*, 1983). Commercial heat treatments (usually at 56°C) to control insects in imported wood packaging have been approved in Japan, China, New Zealand, Australia, Europe, North America and most of South America (Anon., 2005). More information on heat treatments for grain may be found in Chapter 8, this volume.

Heat treatment for structures

The application of dry heat to structures to eliminate pests is an attractive alternative to fumigation, and has been used successfully for decades. Sheppard (1984), Heaps (1988, 1996) and Heaps and Black (1994) recount the measures used in commercial mills and food plants when using heat to control stored-grain pests residing in the infrastructures. More recent work has better defined the treatment parameters (Mahroof *et al.*, 2003; Akdoğan *et al.*, 2005). Detailed information on structural heat treatments for control of storage insects can be found in Chapter 8, this volume.

In addition to stored-product pests, structural heat treatments have been applied to a variety of other structural pests, including roaches (Forbes and Ebeling, 1987), termites (Forbes and Ebeling, 1987; Lewis and Haverty, 1996; Zeichner *et al.*, 1998), ants (Forbes and Ebeling, 1987), and powderpost beetles (Ebeling *et al.*, 1989). Ebeling (1994) reviewed the thermal parameters

and equipment needed to control structural pests. Quarles (1994) noted that thermal treatments were as effective as conventional fumigations for structural pest control and that their costs were decreasing.

High-temperature controlled atmospheres

Another anaerobic procedure combines forced hot air with an oxygen-poor environment, achieved by replacing oxygen with nitrogen or using high concentrations of carbon dioxide. The mechanism of control is to increase respiratory demands for the target pest – as during heating – yet restrict the amount of oxygen available, leading to metabolic arrest and death. Besides exchanging gases, sophisticated instrumentation may be used like those in the forced hot air system.

The greatest advancement and application of high-temperature controlled atmosphere (HTCA) treatments has been against stored-product pests. Many studies on controlled atmosphere (CA) targeting stored-product pests have noted the relationship between temperature and mortality (Harein and Press, 1968; AliNiazee, 1972; Storey, 1975, 1977; Banks and Annis, 1977; Bailey and Banks, 1980; Soderstrom *et al.*, 1986; Delate *et al.*, 1990; Wang *et al.*, 2001). High-temperature controlled atmosphere treatments have now replaced methyl bromide fumigation, particularly in Europe, for pest control of stored food items, spices, grain in silos and ships, furniture and floorboards (Bergwerff and Vroom, 2003).

Studies on HTCA treatments for horticultural crops are of more recent origin. Early experimental units were based on forced hot air design (Gaffney and Armstrong, 1990; Gaffney *et al.*, 1990; Sharp *et al.*, 1991; Neven and Mitcham, 1996). Many HTCA studies for horticultural pests were for control of the light brown apple moth, *Epiphyas postvittana* (Walker) (Lepidoptera: Tortricidae), and other pests of New Zealand apples (Whiting *et al.*, 1991, 1995, 1999a, b; Dentener *et al.*, 1992; Lay-Yee *et al.*, 1997; Chervin *et al.*, 1998; Whiting and Hoy, 1998).

Neven and Mitcham (1996) discussed the development of an HTCA treatment known as the Controlled Atmosphere Temperature Treatment System (CATTS). Applications that show promise include codling moth on fresh sweet cherries (Shellie *et al.*, 2001; Neven, 2005) and nectarines and peaches (Obenland *et al.*, 2005). Neven (2004) provided further discussion of CATTS and other HTCA treatments for fresh commodities. More information on heat with controlled atmospheres may be found in Chapter 10, this volume.

Solar energy

Very little has been done using solar energy to control insects. Although the approach is simple and has potential for long-term storage in rural areas and

developing countries, there are problems with temperature control in terms of the regulation of consistent temperatures. Most work with solar heat treatments of commodities has targeted bruchid pests of seeds using a variety of solar heating methods including plastic bags, corrugated metal and wooden racks (Murdock and Shade, 1991; Kitch *et al.*, 1992; Chinwada and Giga, 1996; Ntougam *et al.*, 1997; Arogba *et al.*, 1998; Songa and Rono, 1998; Ugwu *et al.*, 1999; Chauhan and Ghaffar, 2002).

Solar heating has been used to control other stored product and fruit pests, including: (i) the hide beetle in dried mullet; (ii) the Indianmeal moth in peaches; (iii) the merchant grain beetle in oatmeal (Nakayama *et al.*, 1983); (iv) the larger grain borer in maize cobs (McFarlane, 1989); and (v) nitidulid beetles in figs (Shorey *et al.*, 1989). Solar heating has also been suggested for museum artefacts (Baskin, 2001; Brokerhof, 2003; Pearce, 2003).

Infrared

Infrared is strongly emitted by hot substances and readily absorbed by living tissue, making it a logical choice for thermal treatment. Considerable research has been carried out using infrared heat to control internal pests of grains (rice weevils, lesser grain borers and Angoumois grain moths) using short exposures to temperatures of 56–68°C (Schroeder and Tilton, 1961; Tilton and Schroeder, 1963; Kirkpatrick *et al.*, 1972).

Kirkpatrick and Tilton (1972) measured mortalities of 12 species of stored product beetles in soft winter wheat and obtained > 99.5% mortality for all when treated at 65°C for less than 1 min. Kirkpatrick (1975) treated wheat infested with rice weevils and lesser grain borers in bulk with infrared and obtained > 93% mortality after 24 h exposure to 43.3°C. More recently, Subramanyam (2004) reported mortality from flameless catalytic infrared heaters on adults of the sawtoothed grain beetle, rice weevil and red flour beetle.

Microwave

The microwave (MW) region of the electromagnetic spectrum is from 1 to 100 GHz, between infrared and FM radio, and it is close to radio frequency range (see Fig. 1.2). Microwaves have been applied to a wide range of products, from soil and museum artefacts to fresh fruits. However, the most predominant efforts with current MW technology are in the control of pests of grain and stored products (Nelson 1973, 1996; Roseberg and Bögl, 1987; Nelson *et al.*, 1998; Wang and Tang, 2001).

A number of studies developing MW grain drying systems noted that insect disinfestation was an additional benefit (Hamid and Boulanger, 1970; Boulanger *et al.*, 1971; Tilton and Vardell, 1982a, b). Langlinais (1989)




Fig. 1.2. The electromagnetic spectrum.

demonstrated that the confused flour beetle and flat grain beetle, *Cryptolestes pusillus* (Schnherr) (Coleoptera: Cucujidae), can be economically controlled by MW exposures. Other applications for MW in durable food commodities include treatment for pecan weevil in pecans (Nelson and Payne, 1982), red flour beetle, sawtoothed grain beetle, almond moth and Indianmeal moth in walnuts (Wilkin and Nelson, 1987) and almond moth in sun-dried figs (Baysal *et al.*, 1998).

Investigations into the MW applications of fresh horticultural commodities have been limited. In tests to control the mango weevil, *Cryptorhynchus mangiferae* (F.) (Coleoptera: Curculionidae), Seo *et al.* (1970) observed that MW exposures of 45 s resulted in cooked rind and pulp of treated mangoes. Sharp (1994b) described the equipment used to study the effects of MW on grapefruits infested with the Caribbean fruit fly, *Anastrepha suspense* (Loew) (Diptera: Tephritidae). Sharp *et al.* (1999) examined the effect of MW on mature larvae of the Caribbean fruit fly and concluded that rapid heating imposes serious constraints on the use of heat-induced mortality. Ikediala *et al.* (1999) improved treatment efficacy against the third-instar codling moth by adding 1–2 day cold storage after MW treatments.

MW energy has been explored in the treatment of wood and wood products (Hightower *et al.*, 1974; Burdette *et al.*, 1975; Mashek, 1998) and other non-food items (Reagan *et al.*, 1980; Hall, 1981). Lewis and Haverty (1996) compared MW treatments to five other methods of controlling the drywood termite and obtained 99% mortality 4 weeks after treatment.

To control the Asian longhorned beetle, *Anoplophora glabripennis* (Motschulsky) (Coleoptera: Cerambycidae), Fleming *et al.* (2003) found that MWs heated wood to the controlling temperature of 60°C within 5 min compared to 123 min with conventional heating, and recommended MW treatment to eradicate Asian longhorned beetles in solid wood-packing materials. Philbrick (1984) was concerned about seed viability and slight morphological changes in specimens when MWs were used for herbarium pest control. As with the other applications, MWs work best on dried subjects.

Radio frequency

Radio frequency (RF) waves are at the lower frequency range of the electromagnetic spectrum, with longer wavelength, and the most commonly accepted frequencies used for industrial purposes are 13.56 MHz, 27.12 MHz and 40.68 MHz (Tang *et al.*, 2000). RF energy generates internal heat by resistance from a very rapid change in molecular polarity and migration of charged ions.

The advantages of RF heating are: (i) it is very fast; (ii) it can penetrate deep into the target material because of its long wavelength; (iii) it can produce differential heating between the product and the pest; and (iv) it leaves no toxic residues.

Other than investigations with grain insects (Nelson and Kantack, 1966; Nelson, 1996) and recent progress described later in this volume, RF has not maintained the attention it received in the middle of the last century. Early reviews are provided by Ark and Parry (1940), Webber *et al.* (1946), Frings (1952) and Thomas (1952). Later commentaries are given by Whitney *et al.* (1961) and Watters (1962).

Recent pest control efforts using RF have targeted fresh fruits and nuts (Ikediala *et al.*, 2000; Mitcham *et al.*, 2004; Wang *et al.*, 2005b). Wang *et al.* (2005a) demonstrated the commercial feasibility of RF methodologies to control stored-product pests of walnuts at a walnut-packing house. In Chapters 12 and 13 respectively, this volume, the properties of electromagnetic energy treatments and their potential as commercial treatments will be discussed more thoroughly. A comparison of the heat treatment strategies described above is given in Table 1.3.

Table 1.3. Comparison of heat treatment strategies.

Strategy	First used	Commodity	Advantages	Disadvantages
Hot water	1925	Fruits, bulbs, ornamentals, seeds	Simplest, efficient	Surface heating first; fuel costs
Vapour heat	1913	Fruits, vegetables	Relatively simple	Expensive facilities; surface heating first; slow
Forced hot air	1989	Fruits, vegetables	Product quality retained	Expensive facilities; surface heating first; slow
Dry heat	1792	Structures, grains, fibres, museum artefacts, books	Simple; versatile, known technology	Surface heating first; slow
CATTS	1996	Experimental	Faster than other air methods	Surface heating first; complicated, expensive facilities
Solar	1983	Experimental, structures	Simple, inexpensive	Variable effects
Electromagnetic energy	1927	Experimental, grains, seeds, nuts	Very fast; internal heating first	Expensive facilities; variable effects

1.4 Heat Treatments for Microbial Control

Heat treatments also affect microbial populations on commodities and the incidence of natural decay. Heat treatments are applied to some commodities specifically for decay control, while in other cases the heat treatment is applied for the purpose of insect control and has a secondary effect on decay susceptibility. The secondary effect on decay could be beneficial, resulting in a lower incidence of decay, or detrimental, increasing decay susceptibility. The tolerance of the commodity to the heat treatment has the greatest influence on resultant decay susceptibility. In Chapter 7, this volume, the use of heat treatments for microbial control will be discussed.

1.5 Tolerance of Commodities to Heat Treatments

The tolerance of commodities to heat for insect or decay control must be carefully considered in development of such treatments. If the commodity quality is compromised significantly, the treatment will be unsuccessful. Commodity tolerance to the various methods of heating (water, air, RF, etc.) varies. In some cases, product quality and postharvest life is improved by heat treatment: ripening can be delayed, extending storage life; decay incidence may be reduced.

However, most heat treatments, particularly those designed for control of internal insect pests, cause some detrimental effects on product quality. There are some strategies for increasing the tolerance of commodities. The overall goal is to minimize detrimental effects while maintaining treatment efficacy. See Chapter 4, this volume, for detailed information about the tolerance of commodities to heat treatment and Chapter 11, this volume, for a discussion of induced heat tolerance.

1.6 Conclusions

Insects affect the storage, marketing and trade of food products, resulting in considerable economic loss. Methods to prevent or control insect infestations are needed to maintain food quality and allow free movement of produce. Although chemical fumigation is still the most commonly used control method, environmental and safety issues concerning fumigants provide a strong incentive for the development of alternative, non-chemical methods.

The wide variety of heat treatments currently available or in development show great potential as alternatives, but much work is needed to understand better the biological and physical processes involved. This book provides the foundation for advancing the application of postharvest heat treatments for effective pest control.

1.7 References

- Akdoğan, H., Casada, M.E., Dowdy, A.K. and Subramanyam, Bh. (2005) A novel method for analyzing grain facility heat treatment data. *Journal of Stored Products Research* 41, 175–185.
- AliNiazee, M.T. (1972) Susceptibility of the confused and red flour beetles to anoxia produced by helium and nitrogen at various temperatures. *Journal of Economic Entomology* 65, 60–64.
- Anon. (1990) *Public Law 101-549*. Federal Clean Air Act enacted 15 November 1990. Washington, DC.
- Anon. (1995) Montreal protocol on substances that deplete the ozone layer. *Report of Methyl Bromide Technical Options Committee: 1995 Assessment*. United Nations Environmental Program, Ozone Secretariat, Nairobi, Kenya.
- Anon. (2005) ExO₂ by fumigation in a natural way – wood. ExO₂ (The Netherlands, <http://www.eco2.nl/UK/wood.htm>).
- Ark, P.A. and Parry, W. (1940) Application of high-frequency electrostatic fields in agriculture. *Quarterly Review of Biology* 15, 172–191.
- Armstrong, J.W. (1994a) Commodity resistance to infestation by quarantine pests. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 199–211.
- Armstrong, J.W. (1994b) Heat and cold treatments. In: Paull, R.E. and Armstrong, J.W. (eds) *Insect Pests and Horticultural Products: Treatments and Responses*. CAB International, Wallingford, UK, pp. 103–119.
- Armstrong, J.W., Hansen, J.D., Hu, B.K.S. and Brown, S.A. (1989) High-temperature, forced-air quarantine treatment for papayas infested with tephritid fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology* 82, 1667–1674.
- Arogba, S.S., Ugwu, F.M. and Abu, J.D. (1998) The effects of sun-drying surfaces and packaging materials on the storability of cowpea (*Vigna unguiculata*) seed. *Plant Foods for Human Nutrition* 53, 113–120.
- Back, E.A. and Cotton, R.T. (1924) Relative resistance of the rice weevil, *Sitophilus oryza* L., and the granary weevil, *S. granaries* L., to high and low temperatures. *Journal of Agricultural Research* 28, 1043–1044.
- Bailey, S.W. and Banks, H.J. (1980) A review of recent studies of the effects of controlled atmospheres on stored-product pests. In: Shejbal, J. (ed.) *Controlled Atmosphere Storage of Grains*. Elsevier, Amsterdam, pp. 101–118.
- Baker, A.C. (1939) The basis for treatment of products where fruitflies are involved as a condition for entry into the USA. *USDA Circular* 551, 1–7.
- Baker, A.C. (1952) The vapor-heat process. In: USDA (ed.) *Insects: the Yearbook of Agriculture*. US Government Printing Office, Washington, DC, pp. 401–404.
- Banks, H.J. and Annis, P.C. (1977) Suggested procedures for controlled atmosphere storage of dry grain. *Commonwealth Scientific and Industrial Research Organization, Australia, Division of Entomology Technical Paper* 13, 1–23.
- Baskin, B. (2001) Solar bagging: putting sunlight to work to eliminate insect infestations in mere hours. *Western Association for Art Conservation* 23 (2), 4 pp (<http://palimpsest.stanford.edu/waac/wn/wn23/wn23-2/wn23-207.html>).
- Baysal, T., Ural, A., Çakir, M. and Özen, Ç. (1998) Microwave application for the control of dried fig moth. *Acta Horticulturae* 480, 215–219.
- Bergwerff, F. and Vroom, N. (2003) Exterminating insects and vermin in raw materials. *Grain & Feed Milling Technology* November/December, 28–29.
- Boina, D. and Subramanyam, Bh. (2004) Relative susceptibility of *Tribolium confusum* life stages exposed to elevated temperatures. *Journal of Economic Entomology* 97, 2168–2173.
- Boulanger, R.J., Boerner, W.M. and Hamid, M.A.K. (1971) Microwave and dielectric heating systems. *Milling* 153 (2), 18–21, 24–28.

- Brokerhof, A.W. (2003) The solar tent – cheap and effective pest control in museums. *Australian Institute for the Conservation of Cultural Materials Bulletin* 28, 39–46.
- Burdette, E.C., Hightower, N.C., Burns, C.P. and Cain, F.L. (1975) Microwave energy for wood products insect control. *Proceedings of the Microwave Power Symposium* 10, 276–281.
- Burditt, A.K. (1994) Irradiation. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 101–116.
- Carpenter, A. and Potter, M. (1994) Controlled atmospheres. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 171–198.
- CDFA (California Department of Food and Agriculture) (1998) Cherry fruit fly. In: *Plant Quarantine Manual*. CDFA, Sacramento, California, pp. 305.1–305.3.
- Chauhan, Y.S. and Ghaffar, M.A. (2002) Solar heating of seeds – a low cost method to control bruchid (*Callosobruchus* spp.) attack during storage of pigeonpea. *Journal of Stored Products Research* 38, 87–91.
- Chervin, C.C., Jessup, A., Hamilton, A., Kreidl, S., Kulkarni, S. and Franz, P. (1998) Non-chemical disinfestations: combining the combinations – additive effects of the combination of three postharvest treatments on insect mortality and pome fruit quality. *Acta Horticulturae* 464, 273–278.
- Chew, V. (1994) Statistical methods for quarantine treatment data analysis. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 33–46.
- Chinwada, P. and Giga, D.P. (1996) Sunning as a technique for disinfesting stored beans. *Postharvest Biology and Technology* 9, 335–342.
- Chittenden, F.H. (1897) Some insects injurious to stored grain. *USDA Farmer's Bulletin* 45, 1–24.
- Cohen, M. (1967) Hot-water treatment of plant material. *Ministry of Agriculture, Fisheries and Food Bulletin* 201, 1–39.
- Cotton, R.T. (1963) *Pests of Stored Grain and Grain Products*. Burgess Publishing Co., Minneapolis, Minnesota, 318 pp.
- Couey, H.M. and Hayes, C.F. (1986) Quarantine procedure for Hawaiian papaya using fruit selection and a two-stage hot-water immersion. *Journal of Economic Entomology* 79, 1307–1314.
- Couey, H.M., Armstrong, J.W., Hylin, J.W., Thornburg, W., Nakamura, A.N., Linse, E.S., Ogata, J. and Vetro, R. (1985) Quarantine procedure for Hawaii papaya, using a hot-water treatment and high-temperature, low-dose ethylene dibromide fumigation. *Journal of Economic Entomology* 78, 879–884.
- Davis, J.H. (1933) Radio waves kill insect pests. *Scientific American* May, 272–273.
- Dean, G.A. (1913) Mill and stored-grain insects. *Kansas State Agricultural College, Agricultural Experiment Station Bulletin* No. 189, 63 pp.
- Delate, K.M., Brecht, J.K. and Coffelt, J.A. (1990) Controlled atmosphere treatments for control of sweetpotato weevil (Coleoptera: Curculionidae) in stored tropical sweet potatoes. *Journal of Economic Entomology* 83, 461–465.
- Dentener, P.R., Peetz, S.M. and Birtles, D.B. (1992) Modified atmospheres for the postharvest disinfestations of New Zealand persimmons. *New Zealand Journal of Crop and Horticultural Science* 20, 203–208.
- Dermott, T. and Evans, D.E. (1978) An evaluation of fluidized-bed heating as a means of disinfesting wheat. *Journal of Stored Product Research* 14, 1–12.
- Ebeling, W. (1994) The thermal pest eradication system for structural pest control. *The IPM Practitioner* 16 (2), 1–7.
- Ebeling, W., Forbes, C.F. and Ebeling, S. (1989) Heat treatment for powerpost beetles. *The IPM Practitioner* 11 (9), 1–4.
- Evans, D.E., Thorpe, G.R., and Dermott, T. (1983) The disinfestations of wheat in a continuous-flow fluidized bed. *Journal of Stored Products Research* 19, 125–137.
- Fields, P.G. (1992) The control of stored prod-

- uct insects and mites with extreme temperatures. *Journal of Stored Products Research* 28, 89–118.
- Fields, P.G. and White, N.D.G. (2002) Alternative to methyl bromide treatments for stored-product and quarantine insects. *Annual Review of Entomology* 47, 331–359.
- Fleming, M.R., Hoover, K., Janowiak, J.J., Fang, Y., Wang, X., Liu, W., Wang, Y., Hang, X., Agrawal, D., Mastro, V.C., Lance, D.R., Shield, J.E. and Roy, R. (2003) Microwave irradiation of wood packing material to destroy the Asian longhorned beetle. *Forest Products Journal* 53, 46–52.
- Forbes, C.F. and Ebeling, W. (1987) Update: use of heat for elimination of structural pests. *The IPM Practitioner* 9 (8), 1–5.
- Frings, H. (1952) Factors determining the effects of radio-frequency electromagnetic fields on insects and materials they infest. *Journal of Economic Entomology* 45, 396–408.
- Gaffney, J.J. and Armstrong, J.W. (1990) High-temperature forced-air research facility for heating fruits for insect quarantine treatments. *Journal of Economic Entomology* 83, 1954–1964.
- Gaffney, J.J., Hallman, G.J. and Sharp, J.L. (1990) Vapour heat research unit for insect quarantine treatments. *Journal of Economic Entomology* 83, 1965–1971.
- Gaunce, A.P., Madsen, H.F. and McMullen, R.D. (1981) Fumigation with methyl bromide to kill larvae and eggs of codling moth in Lambert cherries. *Journal of Economic Entomology* 74, 154–157.
- Gould, W.P. (1994) Cold storage. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 119–132.
- Greany, P.D. (1994) Plant host status and natural resistance. In: Paull, R.E. and Armstrong, J.W. (eds) *Insect Pests and Horticultural Products: Treatments and Responses*. CAB International, Wallingford, UK, pp. 37–46.
- Grossman, E.F. (1931) Heat treatment for controlling the insect pest of stored corn. *University of Florida Agricultural Experiment Station Bulletin* 239, 1–24.
- Hadjinicolaou, J. (1931) Effect of certain radio waves on insects affect certain stored products. *Journal of the New York Entomological Society* 39, 145–150.
- Hall, D.W. (1981) Microwave: a method to control herbarium insects. *Taxon* 30, 818–819.
- Hallman, G.J. (1994) Controlled atmospheres. In: Paull, R.E. and Armstrong, J.W. (eds) *Insect Pests and Horticultural Products: Treatments and Responses*. CAB International, Wallingford, UK, pp. 121–136.
- Hallman, G.J. and Armstrong, J.W. (1994) Heated air treatments. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 149–163.
- Hamid, M.A.K. and Boulanger, R.J. (1970) Control of moisture and insect infestation by microwave power. *Milling* 152 (5), 25, 28, 30, 31.
- Hansen, J.D. (2001) Ultrasound treatments to control surface pests of fruit. *Hort-Technology* 11, 186–188.
- Hansen, J.D. and Hara, A.H. (1994) A review of postharvest disinfestations of cut flowers and foliage with special reference to tropics. *Postharvest Biology and Technology* 4, 193–212.
- Harein, P.K. and Press, A.F. (1968) Mortality of stored-peanut insects exposed to mixtures of atmospheric gases at various temperatures. *Journal of Stored Products Research* 4, 77–82.
- Harukawa, C. (1941) Heat as means of controlling Angoumois grain-moth. II. Velocity of the rise of wheat temperature during heating. *Berichte des Ohara Instituts für Landwirtschaftliche Forschungen* 8, 455–464.
- Hawkins, L.A. (1932) Sterilization of citrus fruit by heat. *Texas Citriculture* 9, 7–8, 21–22.
- Headlee, T.J. (1911) Burn the chinch bug in winter quarters. *Kansas State Agricultural College, Experiment Station Circular* No. 19, 8 pp.
- Headlee, T.J. and Burdette, R.C. (1929) Some facts relative to the effect of high frequency radio waves on insect activity.

- Journal of the New York Entomological Society* 37, 59–64.
- Heaps, J.W. (1988) Turn on the heat to control insects. *Dairy and Food Sanitation* 8, 416–418.
- Heaps, J.W. (1996) Heat for stored product insects. *The IPM Practitioner* 18 (5/6), 18–19.
- Heaps, J.W. and Black, T. (1994) Using portable rented electric heaters to generate heat and control stored product insects. *Association of Operative Millers Bulletin* July, 6408–6411.
- Heather, N.W. (1994) Pesticide quarantine treatments. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 89–100.
- Hightower, N.C., Burdette, E.C. and Burns, C.P. (1974) Investigation of the use of microwave energy for weed seed and wood products insect control. *Georgia Institute of Technology Final Technical Report Projects E-230–901*, 1–53.
- Husain, M.A. and Bhasin, H.D. (1921) Preliminary observations on lethal temperatures for the larvae of *Trogoderma khapra*, a pest of stored wheat. *Proceedings of the Fourth Entomological Meeting, Pusa*, February 1921, Calcutta, pp. 240–248.
- Ikediala, J.N., Tang, J., Neven, L.G. and Drake, S.R. (1999) Quarantine treatment of cherries using 915 MHz microwaves: temperature mapping, codling moth mortality and fruit quality. *Postharvest Biology and Technology* 16, 127–137.
- Ikediala, J.N., Tang, J., Drake, S.R. and Neven, L.G. (2000) Dielectric properties of apple cultivars and codling moth larvae. *Transactions of the American Society of Agricultural Engineers* 43, 1175–1184.
- Jang, E.B. and Moffitt, H.R. (1994) Systems approaches to achieving quarantine security. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 225–237.
- Kerbel, E.L., Mitchell, F.G. and Mayer, G. (1987) Effect of postharvest heat treatments for insect control on the quality and market life of avocados. *HortScience* 22, 92–94.
- Kirkpatrick, R.L. (1975) Infrared radiation for control of lesser grain borers and rice weevils in bulk wheat (Coleoptera: Bostrichidae and Curculionidae). *Journal of the Kansas Entomological Society* 48, 100–104.
- Kirkpatrick, R.L. and Tilton, E.W. (1972) Infrared radiation to control adult stored-product Coleoptera. *Journal of the Georgia Entomological Society* 7, 73–75.
- Kirkpatrick, R.L., Brower, J.H. and Tilton, E.W. (1972) A comparison of microwave and infrared radiation to control rice weevil (Coleoptera: Curculionidae) in wheat. *Journal of the Kansas Entomological Society* 45, 434–438.
- Kitch, L.W., Ntoukam, G., Shade, R.E., Wolfson, J.L. and Murdock, L.L. (1992) A solar heater for disinfesting stored cowpeas on subsistence farms. *Journal of Stored Product Research* 28, 261–267.
- Kuznetsova, E.A. (1937) Study of the action of the high frequency field on insects. *Review of Applied Entomology (Series A)* 25, 154.
- Landolt, P.J., Chambers, D.L. and Chew, V. (1984) Alternative to the use of probit-9 mortality as a criterion for quarantine treatments of fruit fly-infested fruit. *Journal of Economic Entomology* 77, 285–287.
- Langlinais, S.J. (1989) Economics of microwave treated rice for controlling weevils. *American Society of Agricultural Engineers Paper* 89-3544, 1–16.
- Latta, R. (1932) The vapour-heat treatment as applied to the control of narcissus pests. *Journal of Economic Entomology* 25, 1020–1026.
- Lay-Yee, M., Whiting, D.C. and Rose, K.J. (1997) Response of 'Royal Gala' and 'Granny Smith' apples to high-temperature controlled atmosphere treatments for control of *Epiphyas postvittana* and *Nysius huttoni*. *Postharvest Biology and Technology* 12, 127–136.
- Lewis, V.R. and Haverty, M.I. (1996) Evaluation of six techniques for control of the western drywood termite (Isoptera:

- Kalotermitidae) in structures. *Journal of Economic Entomology* 89, 922–934.
- Lutz, F.E. (1927) A much abused but still cheerful cricket. *Journal of the New York Entomological Society* 35, 307–308.
- Mackie, D.B. (1931) Heat treatments of California fruits from the standpoint of compatibility of the Florida process. *Monthly Bulletin of the California Department of Agriculture* 20, 211–218.
- MAFF (Ministry of Agriculture, Forestry and Fisheries) – Japan (1950) Plant protection law enforcement regulation. *Ministerial Ordinance No. 73, Annexed Table 1*. MAFF, Tokyo.
- Mahroof, R., Subramanyam, Bh. and Eustace, D. (2003) Temperature and relative humidity profiles during heat treatment of mills and its efficacy against *Tribolium castaneum* (Herbst) life stages. *Journal of Stored Product Research* 39, 555–569.
- Mahroof, R., Subramanyam, Bh. and Flinn, P. (2005) Reproductive performance of *Tribolium castaneum* (Coleoptera: Tenebrionidae) exposed to the minimum heat treatment temperature as pupae and adults. *Journal of Economic Entomology* 98, 626–633.
- Mashek, B.H. (1998) Appropriate technology for drywood termite control. *The IPM Practitioner* 20 (2), 7–8.
- McCoy, C.W., Terranova, A.C., Miller, W.R. and Ismail, M.A. (1994) Vapour heat treatment for the eradication of Fuller rose beetle eggs on grapefruit and its effect on fruit quality. *Proceedings of the Florida Horticultural Society* 107, 235–240.
- McDonald, R.E. and Miller, W.R. (1994) Quality and condition maintenance. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 249–277.
- McFarlane, J.A. (1989) Preliminary experiments on the use of solar cabinets for thermal disinfestations of maize cobs and some observations on heat tolerance in *Prostephanus truncatus* (Horn) (Coleoptera: Bostrichidae). *Tropical Science* 29, 75–89.
- McLaren, G.F., Fraser, J.A. and McDonald, R.M. (1997) The feasibility of hot water disinfestations of summerfruit. *Proceedings of the New Zealand Plant Protection Conference* 50, 425–430.
- McLaren, G.F., Fraser, J.A. and McDonald, R.M. (1999) Non-chemical disinfestations of a quarantine pest on apricots. *Acta Horticulturae* 488, 687–690.
- Milliken, F.B. (1916) Methods of controlling grasshoppers. *Kansas State Agricultural College, Agricultural Experiment Station Bulletin No. 215*, 28 pp.
- Mitcham, E.J., Veltman, R.H., Feng, X., de Castro, E., Johnson, J.A., Simpson, T.L., Biasi, W.V., Wang, S. and Tang, J. (2004) Application of radio frequency treatments to control insects in in-shell walnuts. *Postharvest Biology and Technology* 33, 93–100.
- Moffit, H.R., Drake, S.R., Toba, H.H. and Hartsell, P.L. (1992) Comparative efficacy of methyl bromide against codling moth (Lepidoptera: Tortricidae) larvae in 'Bing' and 'Rainier' cherries and confirmation of efficacy of a quarantine treatment for 'Rainier' cherries. *Journal of Economic Entomology* 85, 1855–1858.
- Moore, D., Lord, J.C. and Smith, S.M. (2000) Pathogens. In: Subramanyam, B. and Hagstrum, D.W. (eds) *Alternatives to Pesticides in Stored-Product IPM*. Kluwer Academic Publishers, Boston, Massachusetts, pp. 193–227.
- Morgan, D.L. and Crocker, R.L. (1986) Survival of weevil-infested live oak acorns treated by heated water. *Texas Agricultural Experiment Station PR-4412* (August), 1–5.
- Mouromtseff, I.E. (1933) Oscillator kills grain weevils in a few seconds. *Electrical World* 102, 667.
- Moy, J.H. (ed.) (1985) *Radiation Disinfestations of Food and Agricultural Products*. University Hawaii Press, Honolulu, Hawaii, 424 pp.
- Murdock, L.L. and Shade, R.E. (1991) Eradication of cowpea weevil (Coleoptera: Bruchidae) in cowpeas by solar heating. *American Entomologist* 37, 228–231.

- Nakayama, T.O.M., Allen, J.M., Cummins, S. and Wang, Y.Y.D. (1983) Disinfestation of dried foods by focused solar energy. *Journal of Food Processing and Preservation* 7, 1-8.
- Nation, J.L. and Burditt, A.K. (1994) Irradiation. In: Paull, R.E. and Armstrong, J.W. (eds) *Insect Pests and Horticultural Products: Treatments and Responses*. CAB International, Wallingford, UK, pp. 85-102.
- Navarro, S. and Calderon, M. (1979) Mode of action of low atmospheric pressures on *Ephestia cautella* (Wlk.) pupae. *Experientia* 35, 620-621.
- Navarro, S., Donahaye, J.E., Dias, R., Azrieli, A., Rindner, M., Phillips, T., Noyes, R., Villers, P., Debruin, T., Truby, R. and Rodriguez, R. (2001) Application of vacuum in a transportable system for insect control. In: Donahaye, E.J., Navarro, S. and Leesch, J.G. (eds) *Proceedings of the International Conference of Controlled Atmospheres and Fumigation in Stored Products*, Fresno, California, 29 October-3 November 2000. Executive Printing Services, Clovis, California, pp. 308-315.
- Nelson, S.O. (1973) Insect-control studies with microwaves and other radiofrequency energy. *Bulletin of the Entomological Society of America* 19, 157-163.
- Nelson, S.O. (1996) Review and assessment of radio-frequency and microwave energy for stored-grain insect control. *Transactions of the American Society of Agricultural Engineers* 39, 1475-1484.
- Nelson, S.O. and Kantack, B.H. (1966) Stored-grain insect control studies with radio-frequency energy. *Journal of Economic Entomology* 59, 588-594.
- Nelson, S.O. and Payne, J.A. (1982) Pecan weevil control by dielectric heating. *Journal of Microwave Power* 17, 51-55.
- Nelson, S.O., Bartley, P.G. and Lawrence, K.C. (1998) RF and microwave dielectric properties of stored-grain insects and their implications for potential insect control. *Transactions of the American Society of Agricultural Engineers* 41, 685-692.
- Neven, L.G. (2004) Hot forced air with controlled atmospheres for disinfestations of fresh commodities. In: Dis, R. and Jain, S.M.S.M. (eds) *Production Practices and Quality Assessment of Food Crops. Vol. 4: Post Harvest Treatments*. Springer, New York, pp. 297-315.
- Neven, L.G. (2005) Combined heat and controlled atmosphere quarantine treatments for control of codling moth in sweet cherries. *Journal of Economic Entomology* 98, 709-715.
- Neven, L.G. and Mitcham, E.J. (1996) CATTs (Controlled Atmosphere/Temperature Treatment System): novel tool for the development of quarantine treatments. *American Entomologist* 42, 56-59.
- Nevo, D. (1996) The desert locust, *Schistocerca gregaria*, and its control in the land of Israel and the near east in antiquity, with some reflections on its appearance in Israel in modern times. *Phytoparasitica* 24, 7-32.
- Ntoukam, G., Kitch, L.W., Shade, R.E. and Murdock, L.L. (1997) A novel method for conserving cowpea germplasm and breeding stocks using solar disinfestations. *Journal of Stored Product Research* 33, 175-179.
- Obenland, D., Neipp, P., Mackey, B. and Neven, L. (2005) Peach and nectarine quality following treatment with high-temperature forced air combined with controlled atmosphere. *HortScience* 40, 1425-1430.
- Oosthuizen, M.J. (1935) The effect of high temperature on the confused flour beetle. *Minnesota Technical Bulletin* 107, 1-45.
- Pearce, A. (2003) Heat eradication of insect infestations: the development of a low cost, solar heated treatment unit. *Australian Institute for the Conservation of Cultural Materials Bulletin* 28, 3-8.
- Pepper, J.H. and Strand, A.L. (1935) Superheating as a control for cereal-mill insects. *Montana State College Agricultural Experiment Station Bulletin* 297, 1-26.
- Philbrick, C.T. (1984) Comments on the use of microwave as a method of herbarium insect control: possible drawbacks. *Taxon* 33, 73-74.
- Prusky, D., Fuchs, Y., Kobilier, I., Roth, I.,

- Weksler, A., Shalom, Y., Fallik, E., Zauberman, G., Pesis, E., Akerman, M., Ykutiely, O., Weisblum, A., Regev, R. and Artes, L. (1999) Effect of hot water brushing, prochloraz treatment and waxing on the incidence of black spot decay caused by *Alternaria alternata* in mango fruits. *Postharvest Biology and Technology* 15, 165–174.
- Quarles, W. (1994) Pest control operators and heat treatment. *The IPM Practitioner* 16 (2), 8.
- Reagan, B.M., Chiao-Cheng, J.-H. and Streit, N.J. (1980) Effects of microwave radiation on the webbing clothes moth, *Tineola bisselliella* (Humm.) and textiles. *Journal of Food Protection* 43, 658–663.
- Riherd, C., Nguyen, R. and Brazzel, J.R. (1994) Pest free areas. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 213–223.
- Robertson, J.L., Preisler, H.K. and Frampton, E.R. (1994a) Statistical concept and minimum threshold concept. In: Paull, R.E. and Armstrong, J.W. (eds) *Insect Pests and Horticultural Products: Treatments and Responses*. CAB International, Wallingford, UK, pp. 47–65.
- Robertson, J.L., Preisler, H.K., Frampton, E.R. and Armstrong, J.W. (1994b) Statistical analyses to estimate efficacy of disinfestations treatments. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 47–65.
- Roseberg, U. and Bögl, W. (1987) Microwave pasteurization, sterilization, blanching, and pest control in the food industry. *Food Technology* June, 92–99.
- Sala, F.J., Burgos, J., Condón, S., Lopez, P. and Raso, J. (1999) Effect of heat and ultrasound on microorganisms and enzymes. In: Gould, G.W. (ed.) *New Methods of Food Preservation*. Chapman & Hall, New York, pp. 176–204.
- Saxena, B.P., Sharma, P.R., Thappa, R.K. and Tikku, K. (1992) Temperature induced sterilization for control of three stored grain beetles. *Journal of Stored Products Research* 28, 67–70.
- Schöller, M. and Flinn, P.W. (2000) Parasites and predators. In: Subramanyam, B.H. and Hagstrum, D.W. (eds) *Alternatives to Pesticides in Stored-Product IPM*. Kluwer Academic Publishers, Boston, Massachusetts, pp. 229–271.
- Schroeder, H.W. and Tilton, E.W. (1961) Infrared radiation for the control of immature insects in kernels of rough rice. *USDA Agricultural Marketing Service* 445, 110.
- Seo, S.T., Chambers, D.L., Komura, M. and Lee, C.Y.L. (1970) Mortality of mango weevils in mangoes treated by dielectric heating. *Journal of Economic Entomology* 63, 1977–1978.
- Seo, S.T., Hu, B.K.S., Komura, M., Lee, C.Y. and Harris, E.J. (1974) *Dacus dorsalis* vapour heat treatment in papayas. *Journal of Economic Entomology* 67, 240–242.
- Sharp, J.L. (1986) Hot-water treatment for control of *Anastrepha suspense* (Diptera: Tephritidae) in mangoes. *Journal of Economic Entomology* 79, 706–708.
- Sharp, J.L. (1994a) Hot water immersion. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 133–147.
- Sharp, J.L. (1994b) Microwaves as a quarantine treatment to disinfest commodities. In: Champ, B.R., Highley, E. and Johnson, G.I. (eds) *Postharvest Handling of Tropical Fruits*. Australian Centre for International Agricultural Research Proceedings No. 56, Canberra, Australia, pp. 362–364.
- Sharp, J.L. and Picho-Martinez, H. (1990) Hot-water quarantine treatment to control fruit flies in mangoes imported into the USA from Peru. *Journal of Economic Entomology* 83, 1940–1943.
- Sharp, J.L., Ouye, M.T., Thalman, R., Hart, W., Ingle, S. and Chew, V. (1988) Submersion of 'Francis' mango in hot water as a quarantine treatment for the West Indian fruit fly and the Caribbean fruit fly (Diptera: Tephritidae) in mangoes. *Journal of Economic Entomology* 81, 1431–1436.
- Sharp, J.L., Ouye, M.T., Hart, W., Ingle, S., Hallman, G., Gould, W. and Chew, V.

- (1989a) Immersion of Florida mangoes in hot water as a quarantine treatment for Caribbean fruit fly (Diptera: Tephritidae). *Journal of Economic Entomology* 82, 186–188.
- Sharp, J.L., Ouye, M.T., Ingle, S.J. and Hart, W.G. (1989b) Hot-water quarantine treatment for mangoes from Mexico infested with Mexican fruit fly and West Indian fruit fly (Diptera: Tephritidae). *Journal of Economic Entomology* 82, 1657–1662.
- Sharp, J.L., Ouye, M.T., Ingle, S.J., Hart, W.G., Enkerlin, H., W.R., Celedonio H., H., Toledo A., J., Stevens, L., Quintero, E., Reyes F., J. and Schwarz, A. (1989c) Hot-water quarantine treatment for mangoes from the state of Chiapas, Mexico, infested with Mediterranean fruit fly and *Anastrepha serpentine* (Wiedemann) (Diptera: Tephritidae). *Journal of Economic Entomology* 82, 1663–1666.
- Sharp, J.L., Gaffney, J.J., Moss, J.I. and Gould, W.P. (1991) Hot-air treatment device for quarantine research. *Journal of Economic Entomology* 84, 520–527.
- Sharp, J.L., Robertson, J.L. and Preisler, H.K. (1999) Mortality of mature third-instar Caribbean fruit fly (Diptera: Tephritidae) exposed to microwave energy. *Canadian Entomologist* 131, 71–77.
- Shellie, K.C., Neven, L.G. and Drake, S.R. (2001) Assessing 'Bing' sweet cherry tolerance to a heated controlled atmosphere for insect pest control. *HortTechnology* 11, 308–311.
- Sheppard, K.O. (1984) Heat sterilization (superheating) as a control for stored-grain pests in a food plant. In: Baur, F.J. (ed.) *Insect Management for Food Storage and Processing*. American Association of Cereal Chemists, St Paul, Minnesota, pp. 194–200.
- Shorey, H.H., Ferguson, L. and Wood, D.L. (1989) Solar heating reduces insect infestations in ripening and drying figs. *HortScience* 24, 443–445.
- Sinclair, W.B. and Lindgren, D.L. (1955) Vapour heat sterilization of California citrus and avocado fruits against fruit-fly insects. *Journal of Economic Entomology* 48, 133–138.
- Soderstrom, E.L., Mackey, B.E. and Brandl, D.G. (1986) Interactive effects of low-oxygen atmospheres, relative humidity, and temperature on mortality of two stored-product moths (Lepidoptera: Pyralidae). *Journal of Economic Entomology* 79, 1303–1306.
- Songa, J.M. and Rono, W. (1998) Indigenous methods for bruchid beetle (Coleoptera: Bruchidae) control in stored beans (*Phaseolus vulgaris* L.). *International Journal of Pest Management* 44, 1–4.
- Stark, J.D. (1994) Chemical fumigants. In: Paull, R.E. and Armstrong, J.W. (eds) *Insect Pests and Horticultural Products: treatments and responses*. CAB International, Wallingford, UK, pp. 69–84.
- Storey, C.L. (1975) Mortality of adult stored-product insects in an atmosphere produced by an exothermic inert atmosphere generator. *Journal of Economic Entomology* 68, 316–318.
- Storey, C.L. (1977) Effect of low oxygen atmospheres on mortality of red and confused flour beetles. *Journal of Economic Entomology* 70, 253–255.
- Stout, O.O. and Roth, H.L. (1983) *International Plant Quarantine Manual*. FAO Plant Production and Protection Paper 50, FAO-UN, Rome, 195 pp.
- Subramanyam, Bh. (2004) Hot technology for killing insects. *Milling Journal First Quarter*, 48–50.
- Tang, J., Ikediala, J.N., Wang, S., Hansen, J.D. and Cavalieri, R.P. (2000). High-temperature, short-time thermal quarantine methods. *Postharvest Biology and Technology* 21, 129–145.
- Thomas, A.M. (1952) Pest control by high-frequency electric fields – critical résumé. *Electrical Research Association Technical Report W/T23*, 7–40.
- Tilton, E.W. and Schroeder, H.W. (1963) Some effects of infrared irradiation on the mortality of immature insects in kernels of rough rice. *Journal of Economic Entomology* 56, 727–730.
- Tilton, E.W. and Vardell, H.H. (1982a) Combination of microwaves and partial vacuum for control of four stored-product insects in stored grain. *Journal of the*

- Georgia Entomological Society 17, 106–112.
- Tilton, E.W. and Vardell, H.H. (1982b) An evaluation of a pilot-plant microwave vacuum drying unit for stored-product insect control. *Journal of the Georgia Entomological Society* 17, 133–138.
- Tsuchiya, T. (1943) Experimental studies on the resistance of the rice weevil, *Calandra oryzae* L., to heat. I. Resistance of adult weevil to heat. *Berichte des Ohara Instituts für Landwirtschaftliche Forschungen* 9, 170–190.
- Ugwu, F.M., Ekwu, F.C. and Abu, J.O. (1999) Effect of different sun-drying surfaces on the functional properties, cooking and insect infestation of cowpea seeds. *Bioresource Technology* 69, 87–90.
- Ulrey, D. (1936) New electronic tubes and new uses. *Physics* 7, 97–105.
- USDA (USA Department of Agriculture) (2004) *USDA–Animal and Plant Health Inspection Service–Plant Protection and Quarantine Manual: Hawaii*. Riverdale, Maryland, 26 pp.
- USDA (USA Department of Agriculture) (2005a) *USDA–Animal and Plant Health Inspection Service–Plant Protection and Quarantine Manual: Regulating the Importation of Fresh Fruits and Vegetables*. Riverdale, Maryland, 306 pp.
- USDA (2005b) *USDA–Animal and Plant Health Inspection Service–Plant Protection and Quarantine Treatment Manual: Nonchemical Treatments*. Riverdale, Maryland, pp. 3.1.1–3.6.10.
- USDA (2005c) *USDA–Animal and Plant Health Inspection Service–Plant Protection and Quarantine Treatment Manual: T106 – Vapour Heat*. Riverdale, Maryland, pp. 5.2.69–5.2.75.
- USEPA (US Environmental Protection Agency) (2001) Protection of stratospheric ozone: process for exempting quarantine and pre-shipment applications of methyl bromide. *Rules and Regulations, Federal Register* 66 (139), 37752–37769.
- Vermeire, L.T., Mitchell, R.B., Fuhlendorf, S.D. and Western D.B. (2004) Selective control of rangeland grasshoppers with prescribed fire. *Journal of Range Management* 57, 29–33.
- Vincent, C., Hallman, G., Panneton, B. and Fleurat-Lessard, F. (2003) Management of agricultural insects with physical control methods. *Annual Review of Entomology* 48, 261–281.
- Walker, G.P., Morse, J.G. and Arpaia, M.L. (1996) Evaluation of a high-pressure washer for postharvest removal of California red scale (Homoptera: Diaspididae) from citrus scale. *Journal of Economic Entomology* 89, 148–155.
- Wang, J.J., Tsai, J.H., Zhao, Z.M. and Li, L.S. (2001) Interactive effects of temperature and controlled atmosphere at biologically relevant levels on development and reproduction of the psocid, *Liposcelis bostrychophila* Badoonel (Psocoptera: Liposcelididae). *International Journal of Pest Management* 47, 55–62.
- Wang, S. and Tang, J. (2001) Radio frequency and microwave alternative treatments for insect control in nuts: a review. *International Agricultural Engineering Journal* 10, 105–120.
- Wang, S., Monzon, M., Tang, J., Johnson, J.A. and Mitcham, E. (2005a) Large-scale radio frequency treatments for insect control in walnuts. *Proceedings of the Tenth Annual International Research Conference on Methyl Bromide Alternatives and Emissions Reduction*, San Diego, California, 31 October–3 November.
- Wang, S., Yue, J., Tang, J. and Chen, B. (2005b) Mathematical modelling of heating uniformity for in-shell walnuts subjected to radio frequency treatments with intermittent stirrings. *Postharvest Biology and Technology* 35, 97–107.
- Watters, F.L. (1962) Control of insects in food-stuffs by high-frequency electric fields. *Proceedings of the Entomological Society of Ontario* 92, 26–32.
- Wearing, C.H., Hansen, J.D., Whyte, C., Miller, C.E. and Brown, J. (2001) The potential for spread of codling moth (Lepidoptera: Tortricidae) via commercial sweet cherry fruit: a critical review and risk assessment. *Crop Protection* 20, 465–488.
- Webber, H.H., Wagner, R.P. and Pearson, A.G. (1946) High-frequency electric fields as

- lethal agents for insects. *Journal of Economic Entomology* 39, 487-498.
- Whiting, D.C. and Hoy, L.E. (1997) High-temperature controlled atmosphere and air treatments to control obscure mealy bug (Hemiptera: Pseudococcidae) on apples. *Journal of Economic Entomology* 90, 546-550.
- Whiting, D.C. and Hoy, L.E. (1998) Effect of temperature establishment time on the mortality of *Epiphyas postvittana* (Lepidoptera: Tortricidae) larvae exposed to a high-temperature controlled atmosphere. *Journal of Economic Entomology* 91, 287-292.
- Whiting, D.C., Foster, S.P. and Maindonald, J.H. (1991) Effects of oxygen, carbon dioxide, and temperature on the mortality responses of *Epiphyas postvittana* (Lepidoptera: Tortricidae). *Journal of Economic Entomology* 84, 1544-1549.
- Whiting, D.C., O'Connor, G.M., van den Heuvel, J. and Maindonald, J.H. (1995) Comparative mortalities of six tortricid (Lepidoptera) species to two high-temperature controlled atmospheres and air. *Journal of Economic Entomology* 88, 1365-1370.
- Whiting, D.C., Hoy, L.E., Maindonald, J.H., Connolly, P.G. and McDonald, R.M. (1998) High-pressure washing treatments to remove obscure mealy bug (Homoptera: Pseudococcidae) and light brown apple moth (Lepidoptera: Tortricidae) from harvested apples. *Journal of Economic Entomology* 91, 1458-1463.
- Whiting, D.C., Jamieson, L.E. and Connolly, P.G. (1999a) Effects of sublethal tebufenozide applications on the mortality responses of *Epiphyas postvittana* (Lepidoptera: Tortricidae) larvae exposed to a high-temperature controlled atmosphere. *Journal of Economic Entomology* 92, 445-452.
- Whiting, D.C., Jamieson, L.E., Spooner, K.J. and Lay-Yee, M. (1999b) Combination high-temperature controlled atmosphere and cold storage as quarantine treatment against *Ctenopseustis obliquana* and *Epiphyas postvittana* on 'Royal Gala' apples. *Postharvest Biology and Technology* 16, 119-126.
- Whitney, W.K., Nelson, S.O. and Walkden, H.H. (1961) Effects of high-frequency electric fields on certain species of stored-grain insects. *USDA Market. Research Report* 455, 1-52.
- Whitney, W.R. (1932) Radiothermy. *General Electric Review* 35, 410-413.
- Wilkin, D.R. and Nelson, G. (1987) Control of insects in confectionery walnuts using microwaves. *British Crop Protection Council Monograph* 37, 247-254.
- Yokoyama, V.Y. (1994) Fumigation. In: Sharp, J.L. and Hallman, G.J. (eds) *Quarantine Treatments for Pests of Food Plants*. Westview Press, Boulder, Colorado, pp. 67-87.
- Yokoyama, V.Y., Miller, G.T. and Hartsell, P.L. (1990) Evaluation of a methyl bromide quarantine treatment to control codling moth (Lepidoptera: Tortricidae) on nectarine cultivars proposed for export to Japan. *Journal of Economic Entomology* 83, 466-471.
- Zeichner, B.C., Hoch, A.L. and Wood, Jr., D.F. (1998) Heat and IPM for cockroach control. *The IPM Practitioner* 20 (2), 1-6.